

ANAPHYLAXIS

HOW TO USE EPIPEN AAIS

If you think someone is have an anaphylactic reaction, give the AAI without delay. It will not harm them.

Always consider anaphylaxis in a food-allergic person even if there are no signs of a rash, hives or swelling.


1. Remove the blue safety cap

Grasp the EpiPen in your dominant hand and remove the blue safety cap by pulling straight up. **Remember: Blue to the Sky, Orange to the Thigh!**




2. Position the orange tip

Hold the EpiPen at 90°, approximately 10cm away from the leg, with the orange tip pointing towards the outer thigh.



3. Administer the EpiPen AAI

Jab the EpiPen firmly into the outer thigh at a right angle. Hold firmly for 3 seconds, before removing and safely discarding.



4. Once the EpiPen AAI has been administered call 999

Ask for an ambulance and say "ana-fill-axis".

5. Lie the person down with legs raised immediately

If the person is not already lying down, they should do so, **with legs raised if possible.**

If breathing is difficult, allow them to sit. If they have vomited or feel sick, gently turn them on their side.




6. If there are no signs of improvement after 5 minutes, use a second EpiPen AAI

The person should remain still and lying down until the ambulance arrives. **Don't try to get up, even if you start to feel better.**

7. Start CPR

If there are no signs of life, **start CPR immediately** until help arrives.



For more information on EpiPen AAI's >>



Sign up to the free expiry alert service and receive reminders by text or email when your EpiPen is about to expire >>

