

ANAPHYLAXIS


HOW TO USE JEXT AAIS

If you think someone is have an anaphylactic reaction, give the AAI without delay. It will not harm them.

Always consider anaphylaxis in a food-allergic person even if there are no signs of a rash, hives or swelling.


1. Hold the Jext AAI in the hand you write with

Hold with your thumb closest to the yellow cap. Pull off the yellow cap with your other hand.



2. Place the black injector tip against the outer thigh

Hold the injector at a right angles (approx. 90°) to the thigh.



3. Push the black tip as hard as you can into the outer thigh

Wait until you hear a 'click' confirming the injection has started, then keep it pushed in. Hold the injector firmly in place against the thigh for 10 seconds (a slow count to 10) then remove. The black tip will extend automatically and hide the needle.



4. Massage the injection area for 10 seconds

5. Once the Jext AAI has been administered call 999

Ask for an ambulance and say "ana-fill-axis".



6. Lie the person down with legs raised immediately

If the person is not already lying down, they should do so, with legs raised if possible.

If breathing is difficult, allow them to sit. If they have vomited or feel sick, gently turn them on their side.




7. If there are no signs of improvement after 5 minutes, use a second Jext AAI

The person should remain still and lying down until the ambulance arrives. Don't try to get up, even if you start to feel better.

8. Start CPR

If there are no signs of life, start CPR immediately until help arrives.



For more information
on Jext AAIs >>



Sign up to the free expiry alert service
and receive reminders by text or email when
your Jext AAI is about to expire >>

